arXiv:1412.0562v2 [math.CV] 15 Dec 2014

# ON REGULARIZATION OF PLURISUBHARMONIC FUNCTIONS NEAR BOUNDARY POINTS

SZYMON PLIŚ

Abstract. We prove in an elementary way that for a Lipschitz domain $D \subset \mathbb{C}^n$, all plurisubharmonic functions on $D$ can be regularized near any boundary point.

## 1. Introduction

Let $D \subset \mathbb{C}^n$. Using a local convolution, for any plurisubharmonic function $u$ on $D$, one can find a sequence $u_k$ of smooth plurisubharmonic functions, which decreases to $u$ on compact subsets of $D$. The purpose of this note is to show that (for regular enough $D$) it is possible to choose such a sequence near any point on the boundary.

A domain $D \subset \mathbb{C}^n$ will be called an $\mathcal{S}$-domain if for any plurisubharmonic function $u$ on $D$, one can find a sequence $u_k$ of smooth plurisubharmonic functions on $D$, which decreases to $u$.[1] Note that pseudoconvex domains, tube domains and Riendhard domains are $\mathcal{S}$-domains (see [F-W]).

**Theorem 1.** *Let $D \subset \mathbb{C}^n$ be a domain with the Lipschitz boundary. Then for any boundary point $P$ there is a neighbourhood $U$ of $P$ such that $D \cap U$ is an $\mathcal{S}$-domain with Lipschitz boundary.*

In section 3, by a slight modification of an example from [F-S], we show the necessity of the assumption on the boundary.

Because of the fact that not every smooth domain is an $\mathcal{S}$-domain (see [F]) we have the following (surprising for the author) corollary:

**Corollary 2.** *Being an $\mathcal{S}$-domain is not a local property of the boundary.*

The author was partially supported by the NCN grant 2011/01/D/ST1/04192.
2010 *Mathematics Subject Classification.* 32U05.
*Key words and phrases.* plurisubharmonic function, regularisation.

[1] Of course this notion does not depend on coordinates so we can define it on manifolds (for example compact manifolds are $\mathcal{S}$-domains). Theorem 1 can be also formulated on manifolds.

## 2. Proof

We need the following Lemma:

**Lemma 3.** *Let $D \subset \mathbb{R}^m$ be an open set. Let $u$ be a subharmonic function on $D$ and let $P \in D$. Let $a, b, R, C > 0$, $B = \{x \in \mathbb{R}^m : |x - P| < R\}$, $K = \{(x', x_m) \in \mathbb{R}^{m-1} \times \mathbb{R} = \mathbb{R}^m : -a < x_m < -b|x'|\}$ and $B + K = \{x + y : x \in B, y \in K\} \subset D$. Assume that for any $x \in B$ and $y \in K$*

$$u(x + y) \leq u(x) + \delta(|y|),$$

*where $\delta : (0, +\infty) \to (0, +\infty)$ is such that $\lim_{t \to 0^+} \delta(t) = 0$. Then $u$ is continuous at $P$.*

*Proof:* Let $(x_n)$ be any sequence in $D$ which converges to $P$. Let $S_n = \{x \in \mathbb{R} : |x - P| = 2|x_n - P|\}$, $A_n = S_n \cap (\{x_n\} + K)$ and $B_n = S_n \setminus A_n$. For $n$ large enough $x_n \in B$ and $|x_n - P| \leq \frac{a}{3}$. Hence there is a constant $\alpha > 0$ (which depends only on $b$) such that $\alpha_n = \sigma(A_n) \geq \alpha\sigma(S_n)$ where $\sigma$ is the standard measure on a sphere. Let $M_n = sup_{S_n} u$. Because $u$ is subharmonic we can estiamate:

$$\begin{aligned} u(P) &\leq \sigma(S_n)^{-1} \int_{A_n} u d\sigma + \sigma(S_n)^{-1} \int_{B_n} u d\sigma \\ &\leq \alpha_n (u(x_n) - \delta(3|x_n - P|)) + (1 - \alpha_n) M_n, \end{aligned}$$

hence

$$u(x_n) \geq u(P) + \frac{1 - \alpha_n}{\alpha_n}(u(P) - M_n) + \delta(3|x_n - P|).$$

Letting $n$ to $\infty$ we get

$$\overline{\lim_{n \to \infty}} u(x_n) \geq u(P).$$

Since $u$ is upper semicontinuous the proof is completed. □

The function $P_D f := \sup\{u \in \mathcal{PSH}(D) : u \leq f\}$, where $D \subset \mathbb{C}^n$ and $f$ is a (real) function on $D$, is called a plurisubharmonic envelope of $f$.

*Proof of Theorem 1:* We use the following notation $\mathbb{C}^n \ni z = (a, x) \in (\mathbb{C}^{n-1} \times \mathbb{R}) \times \mathbb{R}$. We put $B = \{a \in \mathbb{C}^{n-1} \times \mathbb{R} : |a| < 1\}$. After an affine change of coordinates we can assume that there is a constant $C > 0$ and a function $F : B \to [3C, 4C]$ such that:
i) $F(a) - F(b) \leq C|a - b|$ for $a, b \in B$,
ii) $\partial D \cap B \times [-5C, 5C] = \{(a, F(a)) : a \in B\}$ and $P = (0, F(0))$,
iii) $0 \in D$.

Let $U = \{(a,x) \in \mathbb{C}^{n-1} \times \mathbb{R} : |a|^2 + \left(\frac{x}{5C}\right)^2 < 1\}$. We will show that $\Omega = D \cap U$ is an $\mathcal{S}$-domain. Observe that

$$\partial\Omega \cap B \times [0, 5C] = \{(a, \hat{F}(a)) : a \in B\}$$

where $\hat{F} = \max\{F(a), 5C\sqrt{1-|a|^2}\}$. By elementary calculations $\hat{F}(a) - \hat{F}(b) \leq \frac{20}{3}C|a-b|$ for $a, b \in B' = \{a \in \mathbb{C}^{n-1} \times \mathbb{R} : |a| < \frac{4}{5}\}$ and therefore the inequality $\hat{F}(a) - \hat{F}(b) \leq C'|a-b|$ holds in a neighbourhood of $\{\hat{F} = F\} \subset B'$ for some $C' < 7C$.

Let

$$K_\varepsilon = \{(a,x) : -\varepsilon < x < -7C|a|\}.$$

Let

$$\Omega_k = \Omega_k(\varepsilon) = \{z \in \Omega : z + kw \in \Omega \text{ for any } w \in \bar{K}_\varepsilon\}, \text{ for } k = 1, 2.$$

Then there is a compact subset $L_\varepsilon \Subset \Omega_1$ such that

$$\operatorname{dist}(z, \partial\Omega) < \operatorname{dist}(z + w, \partial\Omega)$$

for $z \in \Omega_2 \setminus L_\varepsilon$ and $w \in K_\varepsilon$. Observe that for $\varepsilon$ small enough $\partial D \cap \Omega = \partial D \cap \Omega_2$ and the function $d = -\log(\operatorname{dist}(\cdot, \partial\Omega))$ is plurisubharmonic in a neighbourhood of $cl_\Omega(\Omega \setminus \Omega_2)$. Morover, the function $d' = -\log(\operatorname{dist}(\cdot, \partial U))$ is plurisubharmonic on the whole $\Omega$.

Let $u \in \mathcal{PSH}(\Omega)$ and let $\phi_k$ be a sequence of continuous functions on $\Omega$ which decreases to $u$. We can choose an increasing convex function $p : \mathbb{R} \to \mathbb{R}$ such that for a function $\rho = p \circ d$ we have $\lim_{z \to \partial\Omega} \rho - \phi_1 = +\infty$. Put $\tilde{\phi}_k = \max\{\phi_k, \rho - k\}$. Observe that functions $\hat{\phi}_k = P_\Omega \tilde{\phi}_k$ are plurisubharmonic and they decrease to $u$.

Fix $k$ and $\varepsilon > 0$ such that

$$cl_\Omega(\Omega \setminus \Omega_2) \subset int\{\tilde{\phi}_k = \rho' - k\},$$

where $\rho' = p \circ d'$. On $\Omega_2$ we have

$$\tilde{\phi}_k \geq P_{\Omega_2}\tilde{\phi}_k \geq \hat{\phi}_k \geq \rho' - k$$

and therefore the function $v$ given by

$$v(z) = \begin{cases} P_{\Omega_2}\tilde{\phi}_k(z) & \text{for } z \in \Omega_2 \\ \phi_k(z) = \rho'(z) - k & \text{on } \Omega \setminus \Omega_2, \end{cases}$$

is plurisubharmonic on $\Omega$ and smaller than $\tilde{\phi}_k$. Thus $P_{\Omega_2}\tilde{\phi}_k = \hat{\phi}_k|_{\Omega_2}$. Observe that for $z \in \Omega_2$ and $w \in K_\varepsilon$ we have $\tilde{\phi}_k(z) \geq \tilde{\phi}_k(z+w) - \omega(|w|)$, where $\omega$ is the modulus of continuity of a function $\tilde{\phi}_k|_L$ and $L = L_\varepsilon + K_\varepsilon = \{z + w : z \in L_\varepsilon \text{ and } w \in K_\varepsilon\}$. Therefore, $\hat{\phi}(z) \geq \hat{\phi}(z+w) - \omega(|w|)$. By Lemma 3 the function $\hat{\phi}_k|_{\Omega_2}$ is continuous.

Because any $z \in \Omega$ is in $\Omega_2(\varepsilon)$ for some $\varepsilon$ as above, we obtain that the function $\hat{\phi}_k$ is continuous on $\Omega$. Using the Richberg theorem we can

modify the sequence $\hat{\phi}_k$ to a sequence $u_k$ of smooth plurisubharmonic functions which decreases to $u$. $\square$

The approximation by continuous functions, can be proved in the same way in a much more general situation.

**Theorem 4.** *Let $\boldsymbol{F}$ be a constant coefficient subequetion such that all $\boldsymbol{F}$-subharmonic functions are subharmonic and all convex functions are $\boldsymbol{F}$-subharmonic. Let $D \subset \mathbb{R}^n$ be a domain with Lipschitz boundary. Then for any point $P \in \bar{D}$ there is a neighbourhood $U$ of $P$ such that for any function $u \in \boldsymbol{F}(D \cap U)$ there is a sequence $(u_k) \subset \boldsymbol{F}(D \cap U)$ of continuous functions decreasing to $u$.*

Here we use terminology from [H-L][2].

## 3. Example

Similarly as in Lecture 14 in [F-S] we construct a domain $\Omega \subset \mathbb{C}^n$ and a plurisubharmonic function $u$ on $\Omega$ which can not be regularise. Let $A = \{\frac{1}{k} : k \in \mathbb{N}\}$ and a sequence $(x_k) \subset (0,1) \setminus A$ is such that its limit set is equals $\bar{A}$. Put

$$\lambda(z) = \sum_{k=1}^{\infty} c_k \log|z - x_k| \text{ for } z \in \mathbb{C},$$

where $c_k$ is a such sequence of numbers rapidly decreasing to 0, such that
i) $\lambda$ is a subharmonic function on $\mathbb{C}$ and
ii)$\lambda|_A \geq -\frac{1}{2}$.
For $k \in \mathbb{N}$ let $D_k$ be a disc with center $x_k$ such that $\lambda|_{D_k} < -1$. Now, we can define

$$\Omega = \{(z', z_n) \in \mathbb{C}^{n-1} \times \mathbb{C} : |z'|^2 + |z_n - 1|^2 < 1\} \setminus K,$$

where

$$K = \{(z', z_n) \in \mathbb{C}^{n-1} \times \mathbb{C} : |z'| = |z_n| \text{ and } z_n \notin \cup_{k=1}^{\infty} D_k\},$$

and

$$u(z', z_n) = \begin{cases} -1 & \text{for } z \in \Omega \cap D \\ \max\{\lambda(z_n), -1\} & \text{on } \Omega \setminus D, \end{cases}$$

where $D = \{(z', z_n) \in \mathbb{C}^{n-1} \times \mathbb{C} : |z'| < |z_n|\}$.

[2]See also theorem 2.6 there for elementary properties of **F**-subharmonic functions needed in the proof. For the result of local continuous approximation of $F$-subharmonic functions see [**?**]

Let $U$ be any neighbourhood of 0. We can choose numbers $k \in \mathbb{N}$, $0 < r < \frac{1}{k}$ such that

$$\Omega_U := \{(z', z_n) \in \mathbb{C}^{n-1} \times \mathbb{C} : |z'| \leq \frac{2}{k}, |z_n - \frac{1}{k}| \leq r\} \setminus K \subset \Omega \cap U.$$

Let $y_p$ be subsequence of $x_p$ such that $\lim_{p\to\infty} y_p = \frac{1}{k}$ and for all $p$ we have $|y_p - \frac{1}{k}| < r$. Now, we can repeat the argument from [F-S]. If $u_q$ is a sequence of smooth plurisubharmonic functions decreasing to $u$ on $\Omega_U$, then for $q$ sufficiently large $u_q \leq -\frac{3}{4}$ on the set

$$\{(z', z_n) \in \mathbb{C}^{n-1} \times \mathbb{C} : |z'| = \frac{2}{k}, |z_n - \frac{1}{k}| \leq r\} \subset \partial\Omega_U.$$

By the maximum principle (on sets $\{(z', z_n) \in \mathbb{C}^{n-1} \times \mathbb{C} : |z'| \leq \frac{2}{k}, z_n = y_p\} \subset \mathbb{C}^{n-1\times}\{y_p\}$) we also have $u_q(0, y_p) \leq -\frac{3}{4}$ and by continuity of $u_q$ we get $u_q(0, \frac{1}{k}) \leq -\frac{3}{4} < u(0, \frac{1}{k})$. This is a contradiction.

Note that we have not only just proved that for the above $\Omega$ Theorem 1 does not hold but we also have the following stronger result:

**Proposition 5.** *Let $\Omega$ and $u$ be as above. Then the function $u$ can not be smoothed on $U \cap \Omega$ for any neighbourhood $U$ of $0 \in \partial\Omega$.*

## 4. Questions

In this last section we state some open questions related to the content of the note.

**1.** Is it possible to characterize to characterize $\mathcal{S}$-domains? In view of Corollary 2, even in the class of smooth domains, it seems to be a challenging problem.

**2.** Let $D$ be an $\mathcal{S}$-domain and let $f$ be a continuous function which is bounded from below. Is the plurisubharmonic envelope of $f$ continuous? Assume in addition that $f$ is smooth. What is the optimal regularity of $P_D f$? The author does not know answers even in the case of the ball in $\mathbb{C}^n$. Note that if $D$ is not an $\mathcal{S}$-domain, then there is a smooth function $f$ bounded from below such that $P_D f$ is discontinuous.

**3.** What is the optimal assumption about the regularity of the boundary of $D$ in the Theorem 1? Is it enough to assume that $D = int\bar{D}$?

**4.** Let $M$ be a real (smooth or Lipschitz) hypersurface in $\mathbb{C}^n$. Is it true that for any $P \in M$ there exists a smooth pseudoconvex neighbourhood $U \subset B$ such that $M$ divides $U$ into two $\mathcal{S}$-domains?

## References

[F] J. E. Fornæss, *Plurisubharmonic functions on smooth domains*, Math. Scand. 53 (1983), no. 1, 33-38,

[F-S] J. E. Fornæss, B. Stensønes, *Lectures on counterexamples in several complex variables*, Reprint of the 1987 original. AMS Chelsea Publishing, Providence, RI, 2007,

[F-W] J. E. Fornæss, J. Wiegerinck, *Approximation of plurisubharmonic functions*, Ark. Mat. 27 (1989), no. 2, 257-272,

[H-L] F. R. Harvey, H. B. Lawson Jr., *Dirichlet Duality and the Nonlinear Dirichlet Problem on Riemannian Manifolds*, J. Diff. Geom. **88** (2011), 395-482.

Institute of Mathematics, Cracow University of Technology, Warszawska 24, 31-155 Kraków, Poland

*E-mail address*: splis@pk.edu.pl